

## Balleans of bounded geometry and G-spaces

Igor V. Protasov

Communicated by V. I. Sushchansky

**ABSTRACT.** A ballean (or a coarse structure) is a set endowed with some family of subsets which are called the balls. The properties of the family of balls are postulated in such a way that a ballean can be considered as an asymptotical counterpart of a uniform topological space.

We prove that every ballean of bounded geometry is coarsely equivalent to a ballean on some set  $X$  determined by some group of permutations of  $X$ .

### 1. Ball structures and balleans

A *ball structure* is a triple  $\mathcal{B} = (X, P, B)$ , where  $X, P$  are nonempty sets and, for any  $x \in X$  and  $\alpha \in P$ ,  $B(x, \alpha)$  is a subset of  $X$  which is called a *ball of radius  $\alpha$*  around  $x$ . It is supposed that  $x \in B(x, \alpha)$  for all  $x \in X, \alpha \in P$ . The set  $X$  is called the *support* of  $\mathcal{B}$ ,  $P$  is called the *set of radii*. Given any  $x \in X, A \subseteq X, \alpha \in P$  we put

$$B^*(x, \alpha) = \{y \in X : x \in B(y, \alpha)\}, \quad B(A, \alpha) = \bigcup_{a \in A} B(a, \alpha)$$

A ball structure is called

- *lower symmetric* if, for any  $\alpha, \beta \in P$ , there exist  $\alpha', \beta'$  such that, for every  $x \in X$ ,

$$B^*(x, \alpha') \subseteq B(x, \alpha), \quad B(x, \beta') \subseteq B^*(x, \beta);$$

---

*Thanks to my daughters.*

**2000 Mathematics Subject Classification:** 37B05, 54E15.

**Key words and phrases:** ballean, coarse equivalence, G-space.

- *upper symmetric* if, for any  $\alpha, \beta \in P$ , there exist  $\alpha', \beta'$  such that, for every  $x \in X$ ,

$$B(x, \alpha) \subseteq B^*(x, \alpha'), \quad B^*(x, \beta) \subseteq B(x, \beta');$$

- *lower multiplicative* if, for any  $\alpha, \beta \in P$ , there exists  $\gamma \in P$  such that, for every  $x \in X$ ,

$$B(B(x, \gamma), \gamma) \subseteq B(x, \alpha) \cap B(x, \beta);$$

- *upper multiplicative* if, for any  $\alpha, \beta \in P$ , there exists  $\gamma \in P$  such that, for every  $x \in X$ ,

$$B(B(x, \alpha), \beta) \subseteq B(x, \gamma).$$

Let  $\mathcal{B} = (X, P, B)$  be a lower symmetric and lower multiplicative ball structure. Then the family

$$\left\{ \bigcup_{x \in X} B(x, \alpha) \times B(x, \alpha) : \alpha \in P \right\}$$

is a base of entourages for some (uniquely determined) uniformity on  $X$ . On the other hand, if  $\mathcal{U} \subseteq X \times X$  is a uniformity on  $X$ , then the ball structure  $(X, \mathcal{U}, B)$  is lower symmetric and lower multiplicative, where  $B(x, U) = \{y \in X : (x, y) \in U\}$ . Thus, the lower symmetric and lower multiplicative ball structures can be identified with the uniform topological spaces.

We say that a ball structure  $\mathcal{B}$  is a *ballean* if  $\mathcal{B}$  is upper symmetric and upper multiplicative. In this paper we follow terminology from [6, 7]. A structure on  $X$ , equivalent to a ballean, can also be defined in terminology of entourages. In this case it is called a coarse structure [8] or a uniformly bounded space [5]. For motivations to study balleans see also [1, 2, 4].

## 2. Morphisms

Let  $\mathcal{B}_1 = (X_1, P_1, B_1)$ ,  $\mathcal{B}_2 = (X_2, P_2, B_2)$  be balleans. A mapping  $f : X_1 \rightarrow X_2$  is called a  $\prec$ -*mapping* if, for every  $\alpha \in P_1$ , there exists  $\beta \in P_2$  such that, for every  $x \in X_1$ ,

$$f(B_1(x, \alpha)) \subseteq B_2(f(x), \beta)$$

A bijection  $f : X_1 \rightarrow X_2$  is called an *asymorphism* between  $\mathcal{B}_1$  and  $\mathcal{B}_2$  if  $f$  and  $f^{-1}$  are  $\prec$ -mappings.

Let  $\mathcal{B} = (X, P, B)$  be a ballean,  $S$  be a set. Two mappings  $f, f' : S \rightarrow X$  are called *close* if there exists  $\alpha \in P$  such that  $f'(s) \in B(f(s), \alpha)$  for every  $s \in S$ .

Two ballians  $\mathcal{B}_1 = (X_1, P_1, B_1)$  and  $\mathcal{B}_2 = (X_2, P_2, B_2)$  are called *coarsely equivalent* if there exist the  $\prec$ -mappings  $f_1 : X_1 \rightarrow X_2$ ,  $f_2 : X_2 \rightarrow X_1$  such that  $f_1 \circ f_2, f_2 \circ f_1$  are close to the identity mappings  $id_{X_1}, id_{X_2}$ .

Let  $\mathcal{B} = (X, P, B)$  be a ballean. Every non-empty subset  $Y \subseteq X$  determines the subballean  $\mathcal{B}_Y = (Y, P, B_Y)$ , where  $B_Y(y, \alpha) = B(Y, \alpha) \cap Y$ ,  $y \in Y$ ,  $\alpha \in P$ . A subset  $Y$  is called *large* if there exists  $\gamma \in P$  such that  $B(Y, \gamma) = X$ . If  $Y$  is large, then  $\mathcal{B}_Y$  and  $\mathcal{B}$  are coarsely equivalent. We shall use also the following observations. Two ballians  $\mathcal{B}_1 = (X_1, P_1, B_1)$  and  $\mathcal{B}_2 = (X_2, P_2, B_2)$  are coarsely equivalent if and only if there exist the large subsets  $Y_1 \subseteq X_1, Y_2 \subseteq X_2$  such that the subballians  $\mathcal{B}_{Y_1}$  and  $\mathcal{B}_{Y_2}$  are asyomorphic.

### 3. Density and capacity

Let  $\mathcal{B} = (X, P, B)$  be a ballean,  $Y \subseteq X$ ,  $S \subseteq Y$ ,  $\alpha \in P$ . We say that a subset  $S$  is  $\alpha$ -dense in  $Y$  if  $Y \subseteq B(S, \alpha)$ . An  $\alpha$ -density of  $Y$  is the cardinal

$$den_\alpha(Y) = \min\{|S| : S \text{ is an } \alpha\text{-dense subset of } Y\}.$$

A subset  $S$  of  $X$  is called  $\alpha$ -separated if  $B(x, \alpha) \cap B(y, \alpha) = \emptyset$  for all distinct  $x, y \in S$ . An  $\alpha$ -capacity of  $Y$  is the cardinal

$$cap_\alpha(Y) = \sup\{|S| : S \text{ is an } \alpha\text{-separated subset of } Y\}.$$

Let  $\mathcal{B} = (X, P, B)$  be an arbitrary ballean. Replacing every ball  $B(x, \alpha)$  to  $B'(x, \alpha) = B(x, \alpha) \cap B^*(x, \alpha)$ , we get the asyomorphic ballean  $\mathcal{B}' = (X, P, B')$  with  $(B')^* = B'$ . Thus, in what follows we may suppose that  $B^*(x, \alpha) = B(x, \alpha)$  for all  $x \in X$ ,  $\alpha \in P$ .

**Lemma 1.** *Let  $\mathcal{B} = (X, P, B)$  be a ballean,  $Y \subseteq X$ ,  $\alpha, \beta \in P$  and  $B(B(x, \alpha)) \subseteq B(x, \beta)$  for every  $x \in X$ . Then the following statements hold*

- (i)  $den_\beta(Y) \leq cap_\alpha(Y) \leq den_\alpha(Y)$ ;
- (ii) if  $Z \subseteq X$  and  $Y \subseteq B(Z, \alpha)$ , then  $den_\beta(Y) \leq |Z|$ .

*Proof.* (i) Let  $S$  be an  $\alpha$ -separated subset of  $Y$ ,  $D$  be an  $\alpha$ -dense subset of  $Y$ . Then every ball  $B(x, \alpha)$ ,  $x \in D$  has at most one point of  $S$ . Since  $S \subseteq Y \subseteq \bigcup_{x \in D} B(x, \alpha)$ , we have  $|S| \leq |D|$ , so  $cap_\alpha(Y) \leq den_\alpha(Y)$ .

Let  $S$  be a maximal by inclusion  $\alpha$ -separated subset of  $Y$ . Then every ball  $B(x, \alpha)$ ,  $x \in Y$  meets at least one ball  $B(y, \alpha)$ ,  $y \in S$ . It follows that  $Y \subseteq \bigcup_{x \in S} B(x, \beta)$ , so  $S$  is  $\beta$ -dense in  $Y$  and  $den_\beta(Y) \leq cap_\alpha(Y)$ .

(ii) We put  $Z' = \{z \in Z : B(z, \alpha) \cap Y \neq \emptyset\}$  and, for every  $z \in Z'$ , pick some point  $y_z \in B(z, \alpha) \cap Y$ . Then the subset  $\{y_z : z \in Z'\}$  of  $Y$  is  $\beta$ -dense in  $Y$ , so  $den_\beta(Y) \leq |Z'| \leq |Z|$ .  $\square$

#### 4. Locally finite ballean

A ballean  $\mathcal{B} = (X, P, B)$  is called *locally finite* if every ball  $B(x, \alpha)$ ,  $x \in X$ ,  $\alpha \in P$  is finite.

Let  $\mathcal{B} = (X, P, B)$ ,  $\mathcal{B}' = (X', P', B')$  be ballians,  $f : X \rightarrow X'$  be an injective  $\leftarrow$ -mapping. If  $\mathcal{B}'$  is locally finite then  $\mathcal{B}$  is locally finite. In particular, every ballean asyomorphic to a locally finite ballean is locally finite.

We say that a ballean  $\mathcal{B}$  is *coarsely locally finite* if  $\mathcal{B}$  is coarsely equivalent to some locally finite ballean.

**Proposition 1.** *A ballean  $\mathcal{B} = (X, P, B)$  is coarsely locally finite if and only if there exists  $\beta \in P$  such that  $\beta$ -capacity of every ball  $B(x, \gamma)$ ,  $x \in X$ ,  $\gamma \in P$  is finite.*

*Proof.* Let  $\mathcal{B}' = (X', P', B')$  be a locally finite ballean coarsely equivalent to  $\mathcal{B}$ . Then there exist the large subsets  $Y \subseteq X$ ,  $Y' \subseteq X'$  such that the subballians  $\mathcal{B}_Y$  and  $\mathcal{B}_{Y'}$  are asyomorphic. We choose  $\alpha \in P$  such that  $B(Y, \alpha) = X$  and take an arbitrary  $x \in X$ ,  $\gamma \in P$ . Since  $\mathcal{B}_Y$  is locally finite then the subset  $Z = B(B(x, \gamma), \alpha) \cap Y$  is finite. Since  $B(x, \gamma) \subseteq B(Z, \alpha)$ , by Lemma 1 (ii),  $den_\beta(B(x, \gamma)) \leq |Z|$ . Since  $Z$  is finite, by Lemma 1(i),  $\beta$ -capacity of  $B(x, \gamma)$  is finite.

On the other hand, let  $\beta$ -capacity of every ball  $B(x, \gamma)$  is finite. We choose a maximal by inclusion  $\beta$ -separated subset  $Y$  of  $X$ . Clearly,  $Y$  is large in  $X$ , so  $\mathcal{B}_Y$  is coarsely equivalent to  $\mathcal{B}$ . Since  $cap_\beta B(x, \gamma)$  is finite, then  $B(x, \gamma) \cap Y$  is finite. Hence,  $\mathcal{B}_Y$  is locally finite.  $\square$

Every metric space  $(X, d)$  determines the metric ballean  $\mathcal{B}(X, d) = (X, \mathbb{R}^+, B_d)$ , where  $B_d(x, r) = \{y \in X : d(x, y) \leq r\}$ . For criterion of metrizable of ballians see [7, Theorem 2.1.1]. A metric space is called *proper* if every ball  $B_d(x, r)$  is compact.

**Corollary 1.** *Let  $(X, d)$  be a proper metric space. Then the metric ballean  $\mathcal{B}(X, d)$  is coarsely locally finite.*

*Proof.* It suffices to note that an 1-capacity of every ball in  $(X, d)$  is finite, and apply Proposition 1.  $\square$

## 5. Uniformly locally finite ballean

A ballean  $\mathcal{B} = (X, P, B)$  is called *uniformly locally finite* if there exists a function  $h : P \rightarrow \omega$  such that  $|B(x, \alpha)| \leq h(\alpha)$  for all  $x \in X, \alpha \in P$ .

Let  $\mathcal{B} = (X, P, B), \mathcal{B}' = (X', P', B')$  be ballians,  $f : X \rightarrow X'$  be an injective  $\prec$ -mapping. If  $\mathcal{B}'$  is uniformly locally finite then so is  $\mathcal{B}$ . In particular, every ballean asyomorphic to an uniformly locally finite ballean is uniformly locally finite.

We say that a ballean  $\mathcal{B} = (X, P, B)$  has *bounded geometry* if there exist  $\beta \in P$  and a function  $h : P \rightarrow \omega$  such that  $cap_\beta B(x, \alpha) \leq h(\alpha)$  for all  $x \in X, \alpha \in P$ .

Repeating the arguments proving Proposition 1 we get the following statements.

**Proposition 2.** *A ballean  $\mathcal{B} = (X, P, B)$  has bounded geometry if and only if  $\mathcal{B}$  is coarsely equivalent to some uniformly locally finite ballean.*

**Example 1.** Let  $\Gamma(V, E)$  be a connected graph with the set of vertices  $V$  and the set of edges  $E$ . Given any  $u, v \in V$ , we denote by  $d(u, v)$  the length of a shortest path between  $u$  and  $v$ . Then we get the metric space  $(V, d)$  associated with  $\Gamma(V, E)$  and the metric ballean  $\mathcal{B}(V, d)$ . Clearly,  $\mathcal{B}(V, d)$  is uniformly locally finite if and only if there exists a natural number  $r$  such that  $|B_d(v, 1)| \leq r$  for every  $v \in V$ .

**Example 2.** Let  $G$  be a finitely generated group with the identity  $e$ ,  $F$  be a symmetric ( $F = F^{-1}$ ) set of generators of  $G$  such that  $e \notin F$ . The Cayley graph  $Cay(G, F)$  is a graph with the set of vertices  $G$  and set of edges  $\{\{u, v\} : uv^{-1} \in F\}$ . Let  $d_F$  be a path metric on  $Cay(G, F)$ . Then the metric ballean  $\mathcal{B}(G, d_F)$  is uniformly locally finite.

**Example 3.** Let  $G$  be an arbitrary group,  $\mathcal{F}_e$  the family of all symmetric subsets of  $G$  containing  $e$ . Then we get a ballean  $\mathcal{B}(G) = (G, \mathcal{F}_e, B)$ , where  $B(g, F) = Fg$ . Clearly,  $\mathcal{B}(G)$  is uniformly locally finite and in the case  $G$  is finitely generated,  $\mathcal{B}(G)$  is asyomorphic to the ballean  $\mathcal{B}(G, d_F)$  determined in Example 2.

**Example 4.** Let  $G$  be a group and  $X$  be a  $G$ -space with the action of  $G$  on  $X$  defined by  $(g, x) \mapsto g(x)$ . We denote by  $\mathcal{F}_e$  the family of all finite symmetric subsets of  $G$  containing  $e$ . Then we get the ballean  $\mathcal{B}(G, X) = (X, \mathcal{F}_e, B)$ , where  $B(x, F) = \{g(x) : g \in F\}, x \in X, F \in \mathcal{F}_e$ . Clearly,  $\mathcal{B}(G, X)$  is uniformly locally finite.

**Example 5.** Let  $G$  be a groupoid (=inverse semigroup) of partial bijections of a set  $X$ ,  $\mathcal{F}$  be a family of all finite subsets of  $G$  such that

$F = F^{-1}$  for every  $F \in \mathcal{F}$ . Given any  $x \in X$  and  $F \in \mathcal{F}$ , we put  $B(x, F) = \{x\} \cup \{g(x) : g \in F\}$  and get the uniformly locally finite ballean  $\mathcal{B}(G, X)$ .

**Example 6.** Let  $G$  be a locally compact topological group,  $C$  be the family of all compact symmetric subsets of  $G$  containing  $e$ . Then, by Proposition 5.1, the ballean  $\mathcal{B}(G) = (G, C, B)$ , where  $B(x, C) = Cx$ , is of bounded geometry.

**Remark 1.** Let  $G$  be a locally compact group. Does there exist a discrete group  $D$  such that the ballians  $\mathcal{B}(G)$  and  $\mathcal{B}(D)$  are coarsely equivalent? This is so if  $G$  is Abelian or a connected Lie group.

## 6. $G$ -space realization

Let  $\mathcal{B}, \mathcal{B}'$  be ballians with the same support  $X$ . We write  $\mathcal{B} \prec \mathcal{B}'$  if the identity mapping  $id : X \rightarrow X$  is a  $\prec$ -mapping from  $\mathcal{B}$  to  $\mathcal{B}'$ . If  $\mathcal{B} \prec \mathcal{B}'$  and  $\mathcal{B}' \prec \mathcal{B}$ , we identify  $\mathcal{B}$  and  $\mathcal{B}'$  and write  $\mathcal{B} = \mathcal{B}'$ .

Let  $\mathcal{B}$  be a uniformly locally finite ballean with the support  $X$ . Applying Lemma 4.10 from [8], one can show that there exists a groupoid  $G$  of partial bijections of  $X$  such that  $\mathcal{B} = \mathcal{B}(G, X)$  where  $\mathcal{B}(G, X)$  is a ballean determined in Example 5. Our next result states that instead of the groupoid  $G$  we can take some group of permutations of  $X$ .

**Theorem 1.** *For every uniformly locally finite ballean  $\mathcal{B} = (X, P, B)$ , there exists a group  $G$  of permutations of  $X$  such that  $\mathcal{B} = \mathcal{B}(G, X)$ .*

*Proof.* We fix an arbitrary  $\alpha \in P$  and choose  $\beta \in P$  such that

$$B(B(x, \alpha), \alpha) \subseteq B(x, \beta)$$

for each  $x \in X$ . Then we define the graph  $\Gamma_\beta$  with the set of vertices  $X$  and the set of edges  $E_\beta$  defined by the rule:  $\{x, y\} \in E_\beta$  if and only if  $x \in B(y, \beta)$ . Since  $\mathcal{B}$  is uniformly locally finite, there exists a natural number  $n(\alpha)$  such that the local degree of every vertex of  $\Gamma_\beta$  does not exceed  $n(\alpha)$ . By [3, Corollary 12.2], the chromatic number of  $\Gamma_\beta$  does not exceed  $n(\alpha) + 1$ . It follows that we can partition  $X = X_1 \cup \dots \cup X_{n(\alpha)+1}$  so that any two vertices from  $X_j$  are non-adjacent, in particular, every subset  $X_i$  is  $\alpha$ -separated.

Now we fix  $i \in \{1, \dots, n(\alpha) + 1\}$  and, for every vertex  $x \in X_i$ , enumerate the set  $B(x, \alpha) \setminus \{x\} = \{x(1), \dots, x(n_x)\}$ , where  $n_x \leq n(\alpha)$ . Then we define the set  $S_i(\alpha)$  of  $n(\alpha)$  permutations of  $X$  as follows. For each  $j \in \{1, \dots, n(\alpha)\}$  and  $x \in X_i$ , we put  $\pi_j(x) = x(j)$ ,  $\pi_j(x(j)) = x$  if  $j \leq n_x$ , and  $\pi_j(x) = x$  otherwise. Then we extend  $\pi$  to  $X$  putting

$\pi_j(y) = y$  for all  $y \in X \setminus \bigcup_{x \in X_i} \{x, x(j)\}$ . Since  $X_i$  is  $\alpha$ -separated, this definition is correct. Thus, we get the set  $S_i(\alpha) = \{\pi_1, \dots, \pi_{n(\alpha)}\}$  of permutations of  $X$ . We put  $S(\alpha) = S_1(\alpha) \cup \dots \cup S_{n(\alpha)+1}(\alpha)$  and denote by  $G$  the group of permutations of  $X$  generated by  $\bigcup_{\alpha \in P} S(\alpha)$ .

At last we show that the identity mapping  $id : X \rightarrow X$  is an asymorphism between  $\mathcal{B}$  and the ballean  $\mathcal{B}(G, X) = (X, \mathcal{F}_e, B')$  determined in Example 5.4. Given any  $\alpha \in P$  and  $x \in X$ , we have  $B(x, \alpha) \subseteq B'(x, S_\alpha)$ . On the other hand, let  $F$  be a finite subset of  $G$ ,  $g \in F$ . Then there exists  $\alpha_1, \dots, \alpha_m \in P$  and  $s(\alpha_1) \in S(\alpha_1), \dots, s(\alpha_m) \in S(\alpha_m)$  such that  $g = s(\alpha_m) \dots s(\alpha_1)$ . We choose  $\gamma_g \in P$  such that

$$B(\dots(B(B(x, \alpha_1), \alpha_2), \dots), \alpha_m) \subseteq B(x, \gamma_g)$$

for every  $x \in X$ . Then  $B'(x, \{g\}) \subseteq B(x, \gamma_g)$  for every  $x \in X$ . Since  $F$  is finite, there exists  $\gamma \in P$  such that, for each  $x \in X$ , we have  $B'(x, F) \subseteq B(x, \gamma)$ .  $\square$

Sticking together Proposition 1 and Theorem 1 we get the following statement.

**Theorem 2.** *Every ballean of bounded geometry is coarsely equivalent to some ballean  $\mathcal{B}(G, X)$  of  $G$ -space  $X$ .*

We conclude our paper with two applications of Theorem 1.

**Theorem 3.** *Let  $X$  be a set,  $S_X$  be a group of all permutations of  $X$ . Then  $\mathcal{B}(S_X, X)$  is the strongest uniformly locally finite ballean on  $X$ .*

*Proof.* Let  $\mathcal{B}'$  be a uniformly locally finite ballean on  $X$ . Using Theorem 1, we choose a group  $G$  of permutations of  $X$  such that  $\mathcal{B}' = \mathcal{B}(G, X)$ . Since  $G$  is a subgroup of  $S_X$ , we have  $\mathcal{B}' \prec \mathcal{B}(S_X, X)$ .  $\square$

A ballean  $\mathcal{B} = (X, P, B)$  is called *connected* if, for any  $x, y \in X$ , there exists  $\alpha \in P$  such that  $y \in B(x, \alpha)$ . Clearly, a ballean  $\mathcal{B}(G, X)$  of a  $G$ -space is connected if and only if  $G$  acts transitively on  $X$ .

Let  $\mathcal{B}_1 = (X_1, P_1, B_1)$ ,  $\mathcal{B}_2 = (X_2, P_2, B_2)$  be ballians. A mapping  $f : X_1 \rightarrow X_2$  is called a  $\succ$ -mapping if, for every  $\beta \in P_2$ , there exists  $\alpha \in P_1$  such that  $B_2(f(x), \beta) \subseteq f(B_1(x, \alpha))$  for each  $x \in X_1$ . A bijection  $f : X_1 \rightarrow X_2$  is a  $\succ$ -mapping if and only if  $f^{-1}$  is a  $\prec$ -mapping. Thus,  $\mathcal{B}_1$  and  $\mathcal{B}_2$  are asymorphic if and only if there is a bijection  $f : X_1 \rightarrow X_2$  which is a  $\prec$ -mapping and a  $\succ$ -mapping.

**Theorem 4.** *For every connected uniformly locally finite ballean  $\mathcal{B}$  on a set  $X$ , there exist a group  $G$  of permutations of  $X$  and a surjective mapping  $f : G \rightarrow X$  which is a  $\prec$ -mapping and a  $\succ$ -mapping from  $\mathcal{B}(G)$  to  $\mathcal{B}$ .*

*Proof.* Applying Theorem 1, we identify  $\mathcal{B}$  with  $\mathcal{B}(G, X)$  for some group  $G$  of permutations of  $X$ . Then we fix  $x_0 \in X$  and, for every  $g \in G$ , put  $f(g) = g(x_0)$ . Since  $\mathcal{B}$  is connected,  $(G, X)$  is a transitive  $G$ -space, so  $f$  is surjective. For any finite subset  $F$  of  $G$ , we have  $f(Fg) = Fg(x_0) = F(g(x_0)) = F(f(g))$ . It follows that  $f$  is a  $\prec$ -mapping and a  $\succ$ -mapping.

Let  $(G, X)$  be a transitive  $G$ -space,  $x_0 \in X$ . If  $St(x_0) = \{g \in G : g(x_0) = x_0\}$  is finite, applying Theorem 4, it is easy to show that the ballean  $\mathcal{B}(G)$  and  $\mathcal{B}(G, X)$  are coarsely equivalent.  $\square$

**Remark 2.** Let  $(G, X)$  be a transitive  $G$ -space. How to detect whether the ballean  $\mathcal{B}(G, X)$  is asyomorphic (coarsely equivalent) to the ballean  $\mathcal{B}(H)$  of some group  $H$ ?

### References

- [1] A. Dranishnikov, *Asymptotic topology*, Russian Math. Surveys, **55**(2000), 1085-1129.
- [2] M. Gromov, *Asymptotic invariants for infinite groups*, in Geometric Group Theory, vol. **2**, 1-295, Cambridge University Press, 1993.
- [3] F. Harary, *Graph Theory*, Addison-Wesley Publ. Comp., 1969.
- [4] P. Harpe, *Topics in Geometrical Group Theory*, University Chicago Press, 2000.
- [5] V. Nekrashevych, *Uniformly bounded spaces*, Problems in Algebra, **14**, 47-67, Gomel University Press, 1999.
- [6] I. Protasov, T. Banakh, *Ball Structures and Colorings of Groups and Graphs*, Math. Stud. Monogr. Ser., vol. **11**, VNTL, Lviv, 2003.
- [7] I. Protasov, M. Zarichniy, *General Asymptology*, Math. Stud. Monogr. Ser., vol. **12**, VNTL, Lviv, 2007.
- [8] J. Roe, *Lectures on Coarse Geometry*, University Lecture Series, vol. **31**, American Mathematical Society, Providence, RI, 2003.

### CONTACT INFORMATION

#### I. Protasov

Kyiv Taras Shevchenko Univ. (for Department of Cybernetics), Volodymirska str., 64,  
01033 Kyiv, Ukraine  
*E-Mail:* protasov@unicyb.kiev.ua  
*URL:* <http://do.unicyb.kiev.ua/>

Received by the editors: 23.03.2008  
and in final form 23.03.2008.