

Survey of generalized pregroups and a question of Reinhold Baer

Anthony M. Gaglione, Seymour Lipschutz
and Dennis Spellman

Communicated by I. Ya. Subbotin

ABSTRACT. There has been recent interest in Stallings’ Pregroups. (See [2] and [12].) This paper gives a survey of generalized pregroups. We also answer a question of Reinhold Baer [1] on pregroups and answer a generalization of this question for generalized pregroups.

1. Preliminary results

There has been recent interest in Stallings’ Pregroups. For example:

- [12] Pregroups and the Big Powers Condition: Kvaschuk, Miasnikov, Serbin, Algebra and Logic, Vol. 48, No. 3, 2009
- [2] Geodesic Rewriting Systems and Pregroups, Diekert, Duncan, Miasnikov, 2009, Preprint

First we give some preliminary results.

Let P be a nonempty set with a partial operation, called an “add” by Baer [1] (1950). Formally, a partial operation on P is a mapping $m: D \rightarrow P$ where $D \subseteq P \times P$. If (p, q) belongs to D , we denote $m(p, q)$ by pq and say that pq is defined or exists. (Baer denoted $m(p, q)$ by $p + q$.)

An add P will be called a BS-*pre* or simply a *pre* (term invented by Rimlinger [15]) if it satisfies the following three axioms of Stallings:

2000 MSC: Primary 20E06.

Key words and phrases: Pregroups, Kushner Axiom K. small cancellation.

[P1] (Identity) There exists $1 \in P$ such that for all a , we have $1a$ and $a1$ are defined and $1a = a1 = a$.

[P2] (Inverses) For each $a \in P$, there exists $a^{-1} \in P$ such that aa^{-1} and $a^{-1}a$ are defined, and $aa^{-1} = a^{-1}a = 1$

[P4]=[A] (Weak Associative Law) If ab and bc are defined, then $(ab)c$ is defined if and only if $a(bc)$ is defined, in which case $(ab)c = a(bc)$. (We then say the triple abc is defined.)

Remark 1.1. Stallings also gave the axiom:

[P3] If ab is defined, then $b^{-1}a^{-1}$ is defined and $(ab)^{-1} = b^{-1}a^{-1}$.

However, one can show that [P3] follows from [P1], [P2], and [P4].

It is not difficult to show that: (i) inverses are unique in a pree, (ii) if ab is defined, then $(ab)b^{-1} = a$ and $a^{-1}(ab) = b$.

A sequence $X = [a_1, a_2, \dots, a_n]$ of n elements of P is called a *word with length* $|X| = n$. The word $X = [a_1, a_2, \dots, a_n]$ is said to be *defined* if each pair

$$a_1a_2, a_2a_3, \dots, a_{n-1}a_n$$

is defined. A *triple* in X is a subsequence $a_i a_{i+1} a_{i+2}$.

A product $ab = c$ in a pree may be viewed as a triangle as shown in Fig. 1-1. Bob Gilman [4] noted that the associative law is equivalent to the statement that if three triangles in a pree P fit around a common vertex then the perimeter is also a valid triangle in P . Figure 1-2 illustrates the associative law; that is, the side X is equal to $a(bc)$ and also $(ab)c$.




Fig. 1-1 Product $ab = c$




Fig.1-2 Associative law

Definition 1.2. The *universal group* $G(P)$ of a pree P is the group with presentation $G(P) = gp(P; \text{operation } m)$

That is, P is the set of generators for $G(P)$ and the defining relations of $G(P)$ are of the form $z = xy$ where $m(x, y) = z$.

Definition 1.3. A pree P is said to be *group-embeddable* or simply *embeddable* if P can be embedded in its universal group $G(P)$.

Theorem 1.4. *The question of whether or not a finite pree P embeds in its universal group $G(P)$ is undecidable.*

Bob Gilman [4] noted that this theorem is a special case of a result of Trevor Evans [Embeddable and the word problem] which says that if the embedding problem is solvable for a class of finite partial algebras, then the word problem is solvable for the corresponding class of algebras.

Next follows classical examples of embeddable prees.

Example 1.5. Let K and L be groups with isomorphic subgroups A , pictured in Fig. 1-3. Then the amalgam $P = K \cup_A L$ is a pree which is embeddable in $G(P) = K *_A L$, the free product of K and L with A amalgamated. A typical element w in $G(P)$ is of the form $w = a$ in A or $w = x_1y_1 \cdots x_ny_n$ where x_i and y_i come from different factors in $G(P)$ outside of A .



Fig 1-3



Fig 1-4

Example 1.6. Let K, H, L be groups. Suppose K and H have isomorphic groups A , and suppose H and L have isomorphic groups B , pictured in Fig. 1-4. Then the amalgam $P = K \cup_A H \cup_B L$ is a pree which is embeddable in $G(P) = K *_A H *_B L$ the free product of K, H, L with subgroups A and B amalgamated.

Example 1.7. Let $T = (K_i; A_{rs})$ be a *tree* graph of groups with vertex groups K_i , and with edge groups A_{rs} . Here A_{rs} is a subgroup of vertex groups K_r and K_s . Let $P = \bigcup_i (K_i; A_{rs})$, the amalgam of the groups in T . Then P is a pree which is embeddable in $G(P) = *(K_i; A_{rs})$, the *tree product* of the vertex groups K_i with the subgroups A_{rs} amalgamated.

Example 1.8. Let $G = (K_i; A_{rs})$ be a graph of groups with vertex groups K_i and with edge groups A_{rs} . Again A_{rs} is a subgroup of vertex groups K_r and K_s . Let $P = \bigcup_i (K_i; A_{rs})$. Then P is a pree but P may not be embeddable in $G(P) = *(K_i; A_{rs})$, the free product of groups K_i with the subgroups A_{rs} amalgamated. In fact, there are cases where $G(P) = \{e\}$.

2. Stallings' pregroup

Overall Problem: Find additional axioms so that a pree P is embeddable.

Notation: If X is a set of axioms, then an X -pree will be a pree which also satisfies the axioms in X .

Stallings [16] (1971) invented the name "pregroup" for a pree P and the following axiom:

[P5]=[T1] If ab, bc , and cd are defined, then abc or bcd is defined.

[The reason for the 1 in [T1] is explained in Remark 6.3.]

Theorem 2.1. (Stallings): A pregroup P is embedded in $G(P)$.

[Note: A pregroup P is a T1-pree.]

We quickly outline Stallings' proof of the theorem. A word $w = (x_1, x_2, \dots, x_n)$ is *reduced* if no $x_i x_{i+1}$ is defined. Suppose w is reduced and suppose $x_i a$ and $a^{-1} x_{i+1}$ are defined. Then one can show that

$$w * a = (x_1, x_2, \dots, x_i a, a^{-1} x_{i+1}, \dots, x_n)$$

is also reduced. Stallings called $w * a$ an *interleaving* of w by a .

Define $w \approx v$ if v can be obtained from w by a sequence of interleavings.

Lemma 2.2. $w \approx v$ is an equivalence relation on the set of reduced words.

Lemma 2.3. For any $a \in P$, we define f_a on reduced words by:

$$f_a(x_1, x_2, \dots, x_n) = \begin{cases} (a, x_1, x_2, \dots, x_n) & \text{if } ax_1 \text{ is not defined,} \\ (ax_1, x_2, \dots, x_n) & \text{if } ax_1 \text{ is defined,} \\ & \text{but } ax_1 x_2 \text{ is not defined,} \\ (ax_1 x_2, x_3, \dots, x_n) & \text{if } ax_1 x_2 \text{ is defined.} \end{cases}$$

Lemma 2.4. f_a is a permutation on the equivalence classes of reduced words.

Lemma 2.5. (Main Lemma): If ab is defined then $f_{ab} = f_a f_b$.

The proof of the main lemma consists of the nine possibilities of f_{ab} .

Theorem 2.6. $G(P) = \{\text{permutations } f_a\}$ and P is embedded in $G(P)$ by

$$a \mapsto f_a.$$

Remark 2.7. The pree $P = K \cup_A L$ in Example 1.5 is an example of a pregroup.

3. Baer's question

Reinhold Baer ["Free sums of groups and their generalizations", 1950, [1]] also considered the embedding of prees. In particular, the following appears in his paper:

Postulate XI: (Consists of three parts)

- (a) If ab, bc, cd exist, then $a(bc)$ or $(bc)d$ exist.
- (b) If bc, cd and $a(bc)$ exist, then ab or $(bc)d$ exist.
- (c) If ab, bc and $(bc)d$ exist, then $a(bc)$ or cd exist.

Baer then states:

"In certain instances it is possible to deduce properties (b), (c) from (a); but whether or not this is true in general, the author does not know."

The following theorem (L. and Shi, [14]) answers Baer's question:

Theorem 3.1. *The following conditions on a pree P are equivalent.*

- (i) **[P5] = [T1]:** *If ab, bc, cd are defined, then $a(bc)$ or $(bc)d$ is defined.*
- (ii) **[A1]:** *If $ab, (ab)c, ((ab)c)d$ are defined then bc or cd is defined.*
- (iii) **[A2]:** *If $cd, b(cd), a(b(cd))$ are defined, then ab or bc is defined.*
- (iv) **[A3]:** *If $bc, cd, a(bc)$ are defined, then ab or $(bc)d$ is defined.*
- (v) **[A4]:** *If $ab, bc, (bc)d$ are defined, then $a(bc)$ or cd is defined.*

Note: **[P5] = [T1]** is Baer's (a), **[A3]** is Baer's (b) and **[A4]** is Baer's (c).

Corollary 3.2. *Let P be a pree which satisfies one of the axioms in Theorem 3.1. Then P is embeddable in its universal group $G(P)$.*

4. Kushner's generalization of a pregroup. T2-prees

Note again that $G = K *_A L$ in Example 1.5 is a pregroup since **[P5] = [T1]** does hold in G . However, $G = K *_A H *_B L$ in Example 1.6 is not a pregroup since **[P5] = [T1]** does not hold in G . For example, let $x \in K \setminus A, y \in L \setminus B, a \in A, b \in B$, as pictured in Fig. 4-1. Then $xa \in K, ab \in H$ and $by \in L$ are defined, but xab and aby need not be defined.




Fig.4-1

On the other hand, $G = K *_A H *_B L$ does satisfy the axiom:

[T2] If ab, bc, cd, de are defined, then abc, bcd , or cde is defined. That is, if $X = [a, b, c, d, e]$ is defined, then a triple in X is defined.

Theorem 4.1 (Kushner). *Let P be a T2-pree. Then P is embeddable in $G(P)$.*

We outline the proof of Kushner’s theorem.

Recall that in a pregroup, a reduced word is still reduced under an interleaving. This is not true for a T2-pree. For example, let $x \in K \setminus A, y \in L \setminus B, a \in A, b \in B$, as pictured in Fig.4-1. The word $w = [x, ab, y]$ is reduced in $G = K *_A H *_B L$. But

$$w * a = [xa, a^{-1}(ab), y] = [xa, b, y]$$

is not reduced since by is defined. Thus a reduced word in a T2-pree may not be reduced by an interleaving.

The following definitions are new.

Definition 4.2. The word $w = (x_1, x_2, \dots, x_n)$ is *fully reduced* if w is reduced and w is reduced under any sequence of interleavings.

Definition 4.3. Suppose $w = (x_1, x_2, \dots, x_n)$ is reduced and suppose $x_i = ab$ where $x_{i-1}a$ and bx_{i+1} are defined. Then x_i is said to *split* in w , and w is *reducible* to $v = (x_1, \dots, x_{i-1}a, bx_{i+1}, \dots, x_n)$.

Note first that if w is reducible to v then $|v| < |w|$. Note also that in the above reduced word $w = [x, ab, y]$, the element ab splits in w , and w is reducible to $v = [xa, by]$.

Lemma 4.4. (Main Lemma) *If $w = (x_1, x_2, \dots, x_n)$ is reduced in a T2-pree P , but not fully reduced, then some x_i in w splits.*

That is, w is fully reduced if and only if w is *nonsplitable*.

Define $w \approx v$ if v can be obtained from w by a sequence of interleavings.

Lemma 4.5. *$w \approx v$ is an equivalence relation on the set of fully reduced words.*

If $w = (x_1, x_2, \dots, x_n)$ is fully reduced, then $f_a(w) = f_a(x_1, x_2, \dots, x_n)$ has 5 possible cases (rather the 3 in a pregroup). Thus then following lemma requires 25 cases (not 9).

Lemma 4.6. $f_{ab} = f_a f_b$.

Theorem 4.7. $G(P) = \{\text{permutations } f_a\}$ and P is embedded in $G(P)$ by

$$a \mapsto f_a.$$

5. Baer's question for T2-prees. Open questions for T2-prees

The following theorem generalizes Baer's question for the axiom [T2].

Theorem 5.1 (Gaglione, L, Spellman, 2010). *The following are equivalent in a pree P where a, b, c, d, e are elements in P .*

- 1) [T2] *If ab, bc, cd, de are defined, then $a(bc), b(cd)$, or $c(de)$ is defined.*
- 2) [B1] *If $bc, cd, a(bc), (cd)e$ are defined, then $ab, (bc)d$, or de is defined.*
- 3) [B2] *If $ab, (ab)c, de, c(de)$ are defined, then bc, cd , or $(ab)c(de)$ is defined.*

We Prove Theorem 5.1 in Section 9.

5.1. Transitive order in a pree

The following transitive order relation on a pree P is due to Stallings:

Definition 5.2. Let $L(x) = \{a \in P : ax \text{ is defined}\}$. Put $x \leq y$ if $L(y) \subseteq L(x)$ and $x < y$ if $L(y) \subseteq L(x)$ and $L(y) \neq L(x)$. Also, we let $x \sim y$ if $L(x) = L(y)$.

Example 5.3. Let $P = K \cup_A L$ as in Fig.1-3. Let $x \in K \setminus A$, $y \in K \setminus A$, and $a \in A$. Then $L(x) = K$, $L(y) = K$, $L(a) = P$. Thus, $a < x$ and $a < y$. Also, $x \sim y$.

Theorem 5.4 (Rimlinger, Hoare). *The following conditions on a pree P are equivalent.*

- (i) [P5] = [T1]: *If ab, bc, cd are defined, then $a(bc)$ or $(bc)d$ is defined.*

- (ii) If $x^{-1}a$ and $a^{-1}y$ are defined but $x^{-1}y$ is not defined, then $a < x$ and $a < y$.
- (iii) If $x^{-1}y$ is defined, then $x \leq y$ or $y \leq x$.

Problem (1): Find analogous conditions which are equivalent to [T2].

Theorem 5.5. (Hoare, Chiswell) *The universal group $G(P)$ of a pregroup P admits an integer-valued length function in the sense of Lyndon.*

Problem (2): Prove that an integer-valued length function (in the sense of Lyndon) exists for the universal group $G(P)$ for a T2-pree P .

6. Kushner's axiom K, generalizing [T2]

The proof by Kushner (in his doctoral thesis) that a T2-pree is embeddable was very long and involved (for example, the proof of $f_{ab} = f_a f_b$ required 25 cases instead of 9 cases). Thus the following localization axiom was added in order to shorten the proof:

[K] If ab, bc, cd and $(ab)(cd)$ are defined, then abc or bcd is defined.

Theorem 6.1 (Kushner-L). *Let P be a KT2-pree. Then P is embeddable in $G(P)$.*

After the paper appeared, Hoare independently obtained Kushner's original result with a considerably shorter and less involved proof (by reducing the proof of $f_{ab} = f_a f_b$ to only 9 cases):

Theorem 6.2 (Hoare). *Let P be a T2-pree. Then P is embeddable in $G(P)$.*

Consider the following axioms for $n \geq 1$.

[Tn] If $X = [a_1, a_2, \dots, a_{n+3}]$ is defined, then some triple in X is defined.

That is, if $a_1 a_2, a_2 a_3, \dots, a_{n+2} a_{n+3}$ are defined, then $(a_1 a_2) a_3, (a_2 a_3) a_4, \dots, (a_{n+1} a_{n+2}) a_{n+3}$ is defined.

Remark 6.3. We emphasize that [Tn] holds for a tree pree P in Example 1.7 when the diameter of the tree does not exceed n .

Theorem 6.4 (Kushner-L, 1993). *Let P be a KT3-pree. Then P is embeddable in $G(P)$.*

We note that Theorem 6.4 requires Axiom [K]. The proof of the above theorem again requires:

Lemma 6.5. (Main Lemma) *Let P be a **KT3**-pre. If $w = (x_1, x_2, \dots, x_n)$ is reduced but not fully reduced, then some x_i in w splits.*

Problem (3): Prove that if P is a **T3**-pre, then P is embeddable in $G(P)$.

7. Further generalization

We extend the above Theorem 6.4 to all tree products of groups with finite diameters.

Theorem 7.1. (L) *Let P be a **KTn**-pre. Then P is embeddable in $G(P)$.*

The above theorem requires a generalizing of the notion of a splitting. Specifically:

Definition 7.2. Let $w = (x_1, x_2, \dots, x_n) = (x_1, a_2b_2, a_3b_3, a_4b_4, \dots, a_{n-1}b_{n-1}, x_n)$ where $x_1a_2, b_2a_3, b_3a_4, \dots, b_{n-1}x_n$ are defined. Then we say w is reducible to

$$v = (x_1a_2, b_2a_3, b_3a_4, \dots, b_{n-1}x_n)$$

and the factorization $a_2b_2, a_3b_3, a_4b_4, \dots, a_{n-1}b_{n-1}$ is called a *general splitting* of w .

Remark 7.3. We note that in the above general splitting, $|v| < |w|$.

Example 7.4. Figure 7-1 illustrates a general splitting. Specifically, $w = [x, ab, cd, y]$ need not be reduced where $x \in K_1, y \in K_5, a \in A, b \in B, c \in C, d \in D$. Also, ab need not split and cd need not split. However, xa, bc and dy are defined. Accordingly, $w = [x, ab, cd, y]$ reduces, by a general splitting, to $v = [xa, bc, dy]$.




Fig. 7-1

The following Lemma is essential in the proof of Theorem 7.1.

Lemma 7.5. *Suppose w is reduced but not fully reduced in a \mathbf{KTn} -pree. Then w contains a general splitting.*

We would like to find a theorem which generalizes Bair's question for Axiom $[\mathbf{Tn}]$. Theorem 5.1 answers Baire's question for axiom $[\mathbf{T2}]$. We do have an answer to Baer's question for Axiom $[\mathbf{T6}]$ which we prove in Section 10. Specifically:

Theorem 7.6. *The following axioms, $[\mathbf{T6}]$, $[\mathbf{C6-1}]$, and $[\mathbf{C6-2}]$, are equivalent in a pree P :*

$[\mathbf{T6}]$ *Suppose $X = [a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9]$ is defined, that is, each $a_i a_{i+1}$ is defined. Then a triple in X is defined.*

$[\mathbf{C6-1}]$ *Suppose all the following are defined:*

- (1) $b_2 b_3, b_3 b_4, b_1 (b_2 b_3), (b_3 b_4) b_5,$
- (2) $b_6 b_7, b_7 b_8, b_5 (b_6 b_7), (b_7 b_8) b_9.$

Then one of the following is defined:

$$b_1 b_2, (b_2 b_3) b_4, b_4 b_5, (b_3 b_4) b_5 (b_6 b_7), b_5 b_6, (b_6 b_7) b_8, \text{ or } b_8 b_9.$$

$[\mathbf{C6-2}]$ *Suppose all the following are defined:*

- (1) $b_1 b_2, (b_1 b_2) b_3, b_4 b_5, b_3 (b_4 b_5),$
- (2) $b_5 b_6, (b_5 b_6) b_7, b_8 b_9, b_7 (b_8 b_9).$

Then one of the following is defined:

$$b_2 b_3, (b_1 b_2) b_3 (b_4 b_5), b_3 b_4, (b_4 b_5) b_6, b_6 b_7, (b_5 b_6) b_7 (b_8 b_9), \text{ or } b_7 b_8.$$

Remark 7.7. Note that (2) in both cases $[\mathbf{C6-1}]$ and $[\mathbf{C6-2}]$ can be obtained from (1) by adding 4 to each subscript.

Remark 7.8. The proof of Theorem 7.6 for $[\mathbf{T6}]$ is very similar to the proof of Theorem 5.1 for $[\mathbf{T2}]$ by mainly adding 4 to various subscripts. Likely one can prove an analogous theorem for $[\mathbf{Tm}]$ where $m \equiv 2 \pmod{4}$.

Problem (4): Find a theorem which generalizes Bair's question for axioms $[\mathbf{T3}]$, $[\mathbf{T4}]$ and/or $[\mathbf{T5}]$.

8. Further, further generalizations

Consider Baer's (1953) axioms:

[$S_n, n \geq 4$] Suppose $a_1^{-1}a_2 = b_1, a_2^{-1}a_3 = b_2, \dots, a_{n-1}^{-1}a_n = b_{n-1}, a_n^{-1}a_1 = b_n$ are defined in a pree P . Then at least one of the products $b_i b_{i+1}$ is also defined. (The product may be $b_n b_1$.) In other words, for some $i, a_i^{-1} a_{i+2} \pmod n$ is defined.

Definition 8.1. An S -pree is a pree P which satisfies all axioms S_n for $n \geq 4$.

Axiom S_n is illustrated in Fig. 8-1.




Fig. 8-1

Theorem 8.2. (Baer) Let P be an S -pree. Then P is embeddable in $G(P)$.

Consider two other axioms:

[L] Suppose ab, bc, cd are defined, but $[ab, cd]$ and $[a, bc, d]$ are reduced.

If $(ab)z$ and $z^{-1}(cd)$ are defined, then bz and $z^{-1}c$ are defined.

[M] Equivalent fully reduced words have the same length.

Axiom [M], which we call Baer's axiom, is analogous to his axiom: "Similar irreducible vectors have the same length"

Theorem 8.3. (L, 1996) Let P be a KLM-pree. Then P is embeddable in $G(P)$.

The theorem requires the following proposition which is due to Hoare:

Proposition 8.4 (Hoare). *In a KLM-pree, X is fully reduced if and only if X is nonsplittable.*

Remark 8.5. A KLM-pree includes all tree products of groups, even those without finite diameter.

Theorem 8.6 (Gilman (preprint), Hoare 1998). *Let P be a KL-pree = S_4S_5 -pree. Then P is embeddable in $G(P)$.*

Hoare proved the theorem by showing that axiom [M] follows from [K] and [L].

Gilman proved the theorem using small-cancellation. In particular, Gilman's preprint ["Generalized small cancelation presentations"] indicates an intimate relationship between pregroups and small cancellation theory.

9. Proof of Theorem 5.1

First we restate Theorem 5.1 using different letters for axioms [T2], [B1], and [B2].

Theorem 9.1. *The following are equivalent in a pree P :*

[T2] *If $X = [a_1, a_2, a_3, a_4, a_5]$ is defined, then a triple in X is defined.*

[B1] *If $b_2b_3, b_3b_4, b_1(b_2b_3), (b_3b_4)b_5$ are defined, then one of the following is defined:*

$$b_1b_2, (b_2b_3)b_4 \text{ or } b_4b_5 .$$

[B2] *If $b_1b_2, (b_1b_2)b_3, b_4b_5, b_3(b_4b_5)$ are defined, then one of the following is defined:*

$$b_2b_3, b_3b_4, \text{ or } (b_1b_2)b_3(b_4b_5).$$

Lemma 9.2. [T2] and [B1] are equivalent.

(1) Assume [T2] holds. Suppose the hypothesis of [B1] holds, that is, suppose $b_2b_3, b_3b_4, b_1(b_2b_3), (b_3b_4)b_5$ are defined. Let

$$a_1 = b_1, a_2 = b_2b_3, a_3 = b_3^{-1}, a_4 = b_3b_4, a_5 = b_5.$$

Then the hypothesis of **[T2]** holds, that is, $[a_1, a_2, a_3, a_4, a_5]$ is defined. By **[T2]**, one of the following is defined:

$$a_1a_2a_3 = b_1b_2, \quad a_2a_3a_4 = (b_2b_3)b_4, \quad \text{or} \quad a_3a_4a_5 = b_4b_5.$$

This is the conclusion of **[B1]**. Thus **[T2]** implies **[B1]**.

(2) Assume **[B1]** holds. Suppose the hypothesis of **[T2]** holds, that is, suppose $[a_1, a_2, a_3, a_4, a_5]$ is defined. Let

$$b_1 = a_1, \quad b_2 = a_2a_3, \quad b_3 = a_3^{-1}, \quad b_4 = a_3a_4, \quad b_5 = a_5.$$

Then the hypothesis of **[B1]** holds, that is, $b_2b_3, b_3b_4, b_1(b_2b_3), (b_3b_4)b_5$ are defined. By **[B1]**, one of the following is defined:

$$b_1b_2 = a_1a_2a_3, \quad (b_2b_3)b_4 = a_2a_3a_4, \quad \text{or} \quad b_4b_5 = a_3a_4a_5.$$

This is the conclusion of **[T2]**. Thus **[B1]** implies **[T2]**.

By (1) and (2), **[T2]** and **[B1]** are equivalent in a pree P .

Lemma 9.3. **[T2]** and **[B2]** are equivalent.

(1) Assume **[T2]** holds. Suppose the hypothesis of **[B2]** holds, that is, suppose $b_1b_2, (b_1b_2)b_3, b_4b_5, b_3(b_4b_5)$ are defined. Let

$$a_1 = b_1^{-1}, \quad a_2 = b_1b_2, \quad a_3 = b_3, \quad a_4 = b_4b_5, \quad a_5 = b_5^{-1}.$$

Then the hypothesis of **[T2]** holds, that is, $[a_1, a_2, a_3, a_4, a_5]$ is defined. By **[T2]**, one of the following is defined:

$$a_1a_2a_3 = b_2b_3, \quad a_2a_3a_4 = (b_1b_2)b_3(b_4b_5), \quad \text{or} \quad a_3a_4a_5 = b_3b_4.$$

This is the conclusion of **[B2]**. Thus **[T2]** implies **[B2]**.

(2) Assume **[B2]** holds. Suppose the hypothesis of **[T2]** holds, that is, suppose $[a_1, a_2, a_3, a_4, a_5]$ is defined. Let

$$b_1 = a_1^{-1}, \quad b_2 = a_1a_2, \quad b_3 = a_3, \quad b_4 = a_4a_5, \quad b_5 = a_5^{-1}.$$

Then the hypothesis of **[B2]** holds, that is, $b_1b_2, (b_1b_2)b_3, b_4b_5, b_3(b_4b_5)$ are defined. By **[B2]**, one of the following is defined:

$$b_2b_3 = a_1a_2a_3, \quad b_3b_4 = a_3a_4a_5, \quad \text{or} \quad (b_1b_2)b_3(b_4b_5) = a_2a_3a_4.$$

This is the conclusion of **[T2]**. Thus **[B2]** implies **[T2]**.

By (1) and (2), **[T2]** and **[B2]** are equivalent in a pree P .

Lemma 9.2 and Lemma 9.3 prove Theorem 5.1.

10. Proof of Theorem 7.6.

First we restate Theorem 7.6.

Theorem 10.1. *The following are equivalent in a pree P , where $a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9$ are elements in P .*

[T6] *Suppose $X = [a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9]$ is defined, that is, each $a_i a_{i+1}$ is defined. Then a triple in X is defined.*

[C6-1] *Suppose all the following are defined:*

- (1) $b_2 b_3, b_3 b_4, b_1(b_2 b_3), (b_3 b_4) b_5,$
- (2) $b_6 b_7, b_7 b_8, b_5(b_6 b_7), (b_7 b_8) b_9.$

Then one of the following is defined:

$$b_1 b_2, (b_2 b_3) b_4, b_4 b_5, (b_3 b_4) b_5 (b_6 b_7), b_5 b_6, (b_6 b_7) b_8, \text{ or } b_8 b_9.$$

[C6-2] *Suppose all the following are defined:*

- (1) $b_1 b_2, (b_1 b_2) b_3, b_4 b_5, b_3(b_4 b_5),$
- (2) $b_5 b_6, (b_5 b_6) b_7, b_8 b_9, b_7(b_8 b_9).$

Then one of the following is defined:

$$b_2 b_3, (b_1 b_2) b_3 (b_4 b_5), b_3 b_4, (b_4 b_5) b_6, b_6 b_7, (b_5 b_6 b_7 (b_8 b_9)), \text{ or } b_7 b_8.$$

Remark 10.2. Note that (2) in [C6-1] and (2) in [C6-2] can each be obtained from (1) by adding 4 to each subscript.

Lemma 10.3. *In a pree P , axiom [T6] is equivalent to [C6-1].*

(1) Proof that [T6] implies [C6-1].

Assume [T6] holds. Suppose the hypothesis of [C6-1] holds, that is, the following are defined:

- (1) $b_2 b_3, b_3 b_4, b_1(b_2 b_3), (b_3 b_4) b_5,$
- (2) $b_6 b_7, b_7 b_8, b_5(b_6 b_7), (b_7 b_8) b_9.$

Let

$$\begin{aligned} a_1 &= b_1, & a_2 &= b_2 b_3, & a_3 &= b_3^{-1}, & a_4 &= b_3 b_4, \\ a_5 &= b_5, & a_6 &= b_6 b_7, & a_7 &= b_7^{-1}, & a_8 &= b_7 b_8, & a_9 &= b_9. \end{aligned}$$

Then each $a_i a_{i+1}$ is defined, that is, the hypothesis of [T6] holds. By [T6], one of the following is defined:

$$\begin{aligned} a_1 a_2 a_3 &= b_1 b_2, & a_2 a_3 a_4 &= (b_2 b_3) b_4, & a_3 a_4 a_5 &= b_4 b_5, \\ a_4 a_5 a_6 &= (b_3 b_4) b_5 (b_6 b_7), & a_5 a_6 a_7 &= b_5 b_6, & a_6 a_7 a_8 &= (b_6 b_7) b_8, \\ \text{or } a_7 a_8 a_9 &= b_8 b_9. \end{aligned}$$

This is the conclusion of [C6-1]. Thus [T6] implies [C6-1].

(2) Proof that [C6-1] implies [T6].

Assume [C6-1] holds. Suppose the hypothesis of [T6] holds, that is, suppose $a_1 a_2, a_2 a_3, \dots, a_8 a_9$ are defined. Let

$$\begin{aligned} b_1 &= a_1, & b_2 &= a_2 a_3, & b_3 &= a_3^{-1}, & b_4 &= a_3 a_4, \\ b_5 &= a_5, & b_6 &= a_6 a_7, & b_7 &= a_7^{-1}, & b_8 &= a_7 a_8, & b_9 &= a_9. \end{aligned}$$

Then the hypothesis of [C6-1] holds, that is, the following are defined:

$$\begin{aligned} &b_2 b_3, & b_3 b_4, & b_1 (b_2 b_3), & (b_3 b_4) b_5, & b_6 b_7, \\ &b_7 b_8, & b_5 (b_6 b_7), & (b_7 b_8) b_9. \end{aligned}$$

By [C6-1], one of the following is defined:

$$\begin{aligned} &b_1 b_2 = a_1 a_2 a_3, & (b_2 b_3) b_4 &= a_2 a_3 a_4, & b_4 b_5 &= a_3 a_4 a_5, \\ (b_3 b_4) b_5 (b_6 b_7) &= a_4 a_5 a_6, & b_5 b_6 &= a_5 a_6 a_7, & (b_6 b_7) b_8 &= a_6 a_7 a_8, \\ \text{or } b_8 b_9 &= a_7 a_8 a_9. \end{aligned}$$

This is the conclusion of [T6]. Thus [C6-1] implies [T6].

By (1) and (2), Lemma 10.3 is proved.

Lemma 10.4. *In a pree P , axiom [T6] is equivalent to [C6-2].*

1) Proof that [T6] implies [C6-2]

Assume [T6] holds. Suppose the hypothesis of [C6-2] holds, that is, that the following are defined:

$$b_1 b_2, (b_1 b_2) b_3, b_4 b_5, b_3 (b_4 b_5), b_5 b_6, (b_5 b_6) b_7, b_8 b_9, b_7 (b_8 b_9).$$

Let:

$$\begin{aligned} a_1 &= b_1^{-1}, & a_2 &= b_1 b_2, & a_3 &= b_3, & a_4 &= b_4 b_5, & a_5 &= b_5^{-1}, \\ a_6 &= b_5 b_6, & a_7 &= b_7, & a_8 &= b_8 b_9, & a_9 &= b_9^{-1}. \end{aligned}$$

Then each $a_i a_{i+1}$ is defined, that is, the hypothesis of [T6] holds. By [T6], one of the following is defined:

$$\begin{aligned} a_1 a_2 a_3 &= b_2 b_3, \quad a_2 a_3 a_4 = (b_1 b_2) b_3 (b_4 b_5), \quad a_3 a_4 a_5 = b_3 b_4, \quad a_4 a_5 a_6 = (b_4 b_5) b_6, \\ a_5 a_6 a_7 &= b_6 b_7, \quad a_6 a_7 a_8 = (b_5 b_6) b_7 (b_8 b_9), \quad \text{or } a_7 a_8 a_9 = b_7 b_8. \end{aligned}$$

This is the conclusion of [C6-2]. Thus [T6] implies [C6-2].

(2) Proof that [C6-2] implies [T6].

Assume [C6-2] holds. Suppose the hypothesis of [T6] holds, that is, suppose $a_1 a_2, a_2 a_3, \dots, a_8 a_9$ are defined. Let:

$$\begin{aligned} b_1 &= a_1^{-1}, \quad b_2 = a_1 a_2, \quad b_3 = a_3, \quad b_4 = a_4 a_5, \quad b_5 = a_5^{-1}, \\ b_6 &= a_5 a_6, \quad b_7 = a_7, \quad b_8 = a_8 a_9, \quad b_9 = a_9^{-1}. \end{aligned}$$

Then the hypothesis of [C6-2] holds, that is, the following are defined:

$$b_1 b_2, \quad (b_1 b_2) b_3, \quad b_4 b_5, \quad b_3 (b_4 b_5), \quad b_5 b_6, \quad (b_5 b_6) b_7, \quad b_8 b_9, \quad b_7 (b_8 b_9).$$

By [C6-2], one of the following is defined:

$$\begin{aligned} b_2 b_3 &= a_1 a_2 a_3, \quad (b_1 b_2) b_3 (b_4 b_5) = a_2 a_3 a_4, \quad b_3 b_4 = a_3 a_4 a_5, \quad (b_4 b_5) b_6 = a_4 a_5 a_6, \\ b_6 b_7 &= a_5 a_6 a_7, \quad (b_5 b_6) b_7 (b_8 b_9) = a_6 a_7 a_8, \quad \text{or } b_7 b_8 = a_7 a_8 a_9. \end{aligned}$$

This is the conclusion of [T6]. Thus [C6-2] implies [T6].

By (1) and (2), Lemma 10.4 is proved.

Lemma 10.3 and Lemma 10.4, prove Theorem 7.6.

References

- [1] R. Baer, Free sums of groups and their generalizations II, III, Amer. J. Math. 72 (1950), 625-670.
- [2] Diekert, Duncan, Miasnikov, Geodesic Rewriting Systems and Pregroups, (2009), Preprint
- [3] Evans, Trevor, Embeddable and the word problem, Preprint
- [4] R. H. Gilman, Generalized small cancellation presentations, 1998, preprint.
- [5] A. Gaglione, S. Lipschutz, D. Spellman, Note on a Question of Reinhold Baer on Pregroups II, (2011), Preprint
- [6] A. H. M. Hoare, Pregroups and length functions, Math. Proc. Camb. Phil. Soc. (1988), 104, 21-30.
- [7] _____, On generalizing Stallings' pregroup, J. of Algebra 145 (1992), 113-119
- [8] _____, On generalizing Stallings' pregroup II, (1998), preprint.
- [9] H. Kushner, On prestars and their universal groups, Ph. D. Thesis, Temple University, 1987.

- [10] H. Kushner and S. Lipschutz, A generalization of Stallings' pregroup, *J. of Algebra* 119 (1988), 170-184.
- [11] _____, On embeddable prees, *J. of Algebra* 160 (1993), 1-15.
- [12] Kvaschuk, Miasnikov, Serbin, Pregroups and the Big Powers Condition, *Algebra and Logic* 48, No. 3, (2009),
- [13] S. Lipschutz, Generalizing the Baer-Stallings pregroup, *Contemporary Math.* 169 (1994), 415-430.
- [14] S. Lipschutz and W. Shi, Note on a question of Reinhold Baer on pregroups, *Publ. Inst. Math. (Beograd) (N.S.) B* (2000), 53-58.
- [15] F.S. Rimlinger, Pregroups and Bass-Serre Theory, *Mem. Amer. Math. Soc.* no 361 (American Mathematical Society, 1987).
- [16] J. Stallings, *Group Theory and Three-Dimensional Manifolds*, Yale Monograph 4, (Yale University Press, 1971).

CONTACT INFORMATION

- A. M. Gaglione** Department of Mathematics
U.S. Naval Academy
Annapolis, MD 21402, U.S.A.
E-Mail: amg@usna.edu
URL: <http://www.usna.edu>
- S. Lipschutz** Department of Mathematics
Temple University
Philadelphia, PA 19122, U.S.A.
E-Mail: seymour@temple.edu
- D. Spellman** Department of Statistics
Temple University
Philadelphia, PA 19122, U.S.A.

Received by the editors: 07.02.2012
and in final form 03.04.2012.